

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

1 NAME OF MEDICINE

Water for Injections

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Water for Injections Freeflex 100 mL, 250 mL, 500 mL or 1000 mL contains 100% v/v water for injections

3 PHARMACEUTICAL FORM

Injection, intravenous infusion

Water for Injections Freeflex is a sterile, clear, colourless, particle-free, odourless and tasteless liquid with a pH of 5.6-7.7. It contains no anti-microbial agents.

4 CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATIONS

Water for Injections is used to dissolve or dilute substances or preparations for parenteral administration.

Water for Injections may also be used as an irrigating solution for small wounds or during minor surgical procedures.

4.2 DOSE AND METHOD OF ADMINISTRATION

For dissolving or diluting agents for parenteral administration

The dosage for Water for Injections is that required to dissolve or dilute other agents. Aseptic technique must be used when preparing solutions for parenteral administration. Check the Product Information of any substance, preparation or drug before use to ensure appropriate solubility, dilution or compatibility with other additives.

Solutions prepared with Water for Injections may be administered intravenously, intramuscularly or subcutaneously using strict aseptic technique. Care should be exercised that all solutions prepared with Water for Injections are isotonic before use (See Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE). Water for Injections is to be used for one patient on one occasion only. Any residue should be discarded. It does not contain antimicrobials. Care should be taken with intravenous administration and injection technique to avoid injection site reactions and infections.

Usually solutions are prepared immediately before use. The Product Information of substances or drugs to be dissolved or diluted must be consulted to ascertain the maximum time between aseptic preparation and use of the solution.

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

For irrigation

Before using Water for Injections to irrigate small wounds, or during minor surgical procedures, inspect the contents to ensure that there has been no discolouration. Water for Injections is a sterile product and when used for irrigation, strict aseptic technique should be observed at all times. Water for Injections is for use for a single patient on a single occasion. Any residue remaining should be discarded.

4.3 CONTRAINDICATIONS

Water for Injections is hypotonic causing haemolysis if it is injected alone. It is contraindicated for intravenous administration if it is not adjusted to isotonicity by the addition of suitable solutes.

The use of Water for Injection as irrigation during a major surgical procedure, or in a procedure where significant amounts may be absorbed or enter the circulation, is contraindicated.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Do not use Water for Injections unless it is clear and the seal is intact.

Check the Product Information of any substance, preparation or drug before use to ensure appropriate solubility, dilution or compatibility with other additives.

Before intravenous administration of a solution prepared with Water for Injections, ensure that the resultant solution is isotonic with blood. Entry of water or hypotonic solution into the systemic circulation may cause haemolysis. Given that there is a possibility of systemic absorption of irrigation solutions, the same precautions apply.

Tissue damage may result from irrigation with large volumes or under pressure: see Section 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS).

Use in the elderly

No data available

Paediatric Use

No data available

Effects on laboratory tests

No data available

4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTION

Not applicable

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

4.6 FERTILITY, PREGNANCY AND LACTATION

Effects on fertility

No data available.

Use in Pregnancy (Category A)

Check the Product Information document of the drug to be dissolved or diluted to ensure that it is safe to use during pregnancy.

Use in Lactation

Water for Injections can be administered to women who are breastfeeding. Check the Product Information document of the drug to be dissolved or diluted to ensure that it is safe to use during lactation.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

The effects of this medicine on a person's ability to drive and use machines were not assessed as part of its registration.

4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

Haemolysis and hyponatraemia have been reported after irrigation during urological procedure. There should be no adverse reaction to Water for Injections if used as indicated to dissolve compatible substances to form an isotonic solution prior to injection. Injection of Water for Injections without the addition of solute may result in cell damage due to hypotonic effects. (See 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE). Haemolysis may lead to renal tubular obstruction. Expansion of intravascular fluid, through intravenous infusion, or systemic absorption of irrigation solutions, may result in electrolyte disturbances including hyponatraemia, and cardiovascular/pulmonary disorders due to oedema.

The Product Information of any drug or substance used with Water for Injections must be consulted before use.

Intravenous administration of solutions may cause local reactions including pain, vein irritation, and thrombophlebitis. Extravasation of solution may cause tissue injury.

Displaced catheters or drainage tubes can lead to irrigation or infiltration of unintended structures or cavities. Excessive volumes or pressure during irrigation of closed cavities may result in distension or disruption of tissues. Inadvertent contamination from careless techniques may transmit infection.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at www.tga.gov.au/reporting-problems.

4.9 OVERDOSE

If Water for Injections is inadvertently injected without first ensuring isotonicity, the hypotonic effects may include local cell damage or haemolysis.

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

Overdose using isotonic intravenous solutions prepared using Water for Injections or during irrigation, may cause fluid overload and electrolyte disturbances. See Section 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS).

Infusion or irrigation should be ceased and the patient assessed and treated appropriately.

For information on the management of overdose, contact the Poison Information Centre on 131126 (Australia) or 0800 764 766 (New Zealand).

5 PHARMACOLOGICAL PROPERTIES

5.1 PHARMACODYNAMIC PROPERTIES

Mechanism of action

Water is the main constituent of the body fluids. Body weight is approximately 60% of water distributed in intracellular, interstitial and vascular compartments. The water content in the intracellular fluid, ie. the water inside the cells, is about 40 to 45 % of body weight. Water moves freely between these compartments. Thus, pharmacological action of the Water for Injection is as a vehicle for substances in maintaining the isotonicity across these compartments.

Clinical trials

No data available.

5.2 PHARMACOKINETIC PROPERTIES

As Water for Injections is solute-free with osmolarity of zero (a hypotonic solution), its entry into the systemic circulation will result in a dilution of the electrolytes in the extracellular fluid leading to the movement of water into the red blood cells causing haemolysis. Thus, Water for Injections should not be injected without adjusting it to isotonicity by the addition of suitable solute.

5.3 PRECLINICAL SAFETY DATA

Genotoxicity

Water is the main constituent of the body fluids and is not known as a mutagen.

Carcinogenicity

Water is the main constituent of the body fluids and is not known as a carcinogen.

6 PHARMACEUTICAL PARTICULARS

6.1 LIST OF EXCIPIENTS

Not applicable

6.2 INCOMPATIBILITIES

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

However, additives may be incompatible with Water for Injections Freeflex (see Section 4.2 DOSE AND METHOD OF ADMINISTRATION and Section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE).

6.3 SHELF LIFE

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 SPECIAL PRECAUTIONS FOR STORAGE

Store below 25°C.

6.5 NATURE AND CONTENTS OF CONTAINER

Water for Injections Freeflex is supplied in freeflex bags – polyolefin.

Water for Injections Freeflex 100 mL AUST R 144795 – available in packs of 40, 50, 55 and 60 bags

Water for Injections Freeflex 250 mL AUST R 144808 – available in packs of 20, 30, 35 and 40 bags

Water for Injections Freeflex 500 mL AUST R 144809 – available in packs of 20 bags

Water for Injections Freeflex 1000 mL AUST R 144814 – available in packs of 10 bags.

Not all pack sizes are marketed

6.6 SPECIAL PRECAUTIONS FOR DISPOSAL

In Australia, any unused medicine or waste material should be disposed of in accordance with local requirements.

6.7 PHYSICOCHEMICAL PROPERTIES

Chemical structure

Chemical formula is H₂O

Molecular Weight

18.02

CAS number

The CAS number for water is 7732-18-5

7 MEDICINE SCHEDULE (POISONS STANDARD)

Australia: Not scheduled

New Zealand: General sales Medicine

AUSTRALIAN PRODUCT INFORMATION

WATER FOR INJECTIONS FREEFLEX (WATER FOR INJECTION)

8 SPONSOR

Fresenius Kabi Australia Pty Limited
Level 2, 2 Woodland Way
Mount Kuring-gai, NSW 2080
Australia
Telephone: (02) 9391 5555

Fresenius Kabi New Zealand Limited
60 Pavilion Drive
Airport Oaks, Auckland 2022
New Zealand
Freecall: 0800 144 892

9 DATE OF FIRST APPROVAL

11 August 2005

10 DATE OF REVISION

13 December 2018

SUMMARY TABLE OF CHANGES

Section Changed	Summary of new information
All	New format