



Patient reminder card

This card outlines the important safety information that you need to be aware of prior to and throughout treatment with Idacio.

Fresenius Kabi Canada Ltd.

165 Galaxy Blvd, Suite 100,
Toronto, ON M9W 0C8

Medical information contact number:

1-877-821-7724

November 30, 2022

FKC-4-0023-00



you/your child more likely to get infections or make any infection that you may have worse. This includes infections such as colds or more serious infections like tuberculosis.

Cancer

If you/your child take Idacio, the risk of getting certain types of cancer may increase.

Nervous system problems

Some people have developed new or worsening nervous system problems with Idacio. This includes multiple sclerosis. Please read the Idacio package leaflet for more information. These are not all of the possible side effects that may occur during treatment with Idacio.

Call your/your child's doctor or get medical care right away if you/your child have any of the following symptoms of these possible serious side effects:

Infections

- Fever, chills, unusual sweating, feeling unwell or more tired than normal, feeling or being sick (such as nausea or vomiting), diarrhea, stomach pain, loss of appetite or weight loss, cough or coughing up blood or mucus, shortness of breath, problems urinating, skin sores, wounds, muscle aches, dental problems.

Cancer

- Night sweats, swelling of the lymph nodes (swollen glands) in the neck, armpits, groin or other areas, weight loss, new skin lesions or change in skin lesions (such as moles or freckles) you/your child already have, severe itchiness that cannot be explained.

Nervous system problems

- Numbness or tingling, vision changes, muscle weakness, unexplained dizziness.
- Tell your/your child's doctor about any unusual symptoms that you/your child may have during treatment with Idacio. These are not all of the possible symptoms of these side effects.

Information for you/your child and healthcare professionals involved in medical care or treatment.

Tuberculosis (TB) tests and treatment.

Mark this box if you/your child have ever been tested for TB:

YES (Check with the doctor if you do not know)

Mark this box if you/your child have ever had any test that was positive for TB:

YES (Check with the doctor if you do not know)

Mark this box if you/your child have ever taken any pills to treat or prevent TB:

YES (Check with the doctor if you do not know)

Patient name: _____

Doctor's name (who prescribed Idacio): _____

Doctor's phone number: _____

Date of first Idacio injection: _____

Idacio LOT number: _____

Dose of Idacio injection: _____

Date of last Idacio injection (if no longer taking Idacio): _____

Please fill out information for you/your child and healthcare professionals involved in your/your child's medical care or treatment.

Notes (comments or questions for your/your child's doctor): _____

This card contains select important safety information that you need to know before and during your/your child's treatment with Idacio.

- Keep this card with you at all times and for 70 days after your/your child's last injection of Idacio.
- Show this card to any doctor or healthcare professional that you/your child see.
- Record information about any tuberculosis tests or treatment you/your child have had on the back of this card.

Introduction

Idacio is a medicine that is intended to treat certain diseases that affect a part of the immune system. While Idacio can be effective in treating these diseases, some people can have one or more side effects. It is important to talk to your/your child's doctor about the possible benefits and possible side effects of taking Idacio. These can be different for each person.

- The purpose of this card is to tell you about some of the possible side effects of Idacio.
- Some of the serious side effects that could occur include infections, cancer, and nervous system problems.
- These are not all of the possible side effects of Idacio.

Before Idacio treatment

Tell your/your child's doctor about any health problems you/your child have and any medicines you/your child take. This will help you and your/your child's doctor decide if Idacio is right for you/your child.

Tell your/your child's doctor if you/your child:

- Have an infection or have symptoms of an infection (such as fever, wounds, feeling tired, dental problems);
- Have tuberculosis or have been in close contact with someone with tuberculosis;
- Have or have had cancer;
- Have any numbness or tingling or have a problem that affects your/your child's nervous system, such as multiple sclerosis.

Your/your child's doctor should check you/your child for signs and symptoms of tuberculosis before starting Idacio. You/your child may need to be treated for tuberculosis before you/your child start taking Idacio.

Vaccinations

- You/your child may receive vaccinations except for live vaccines.
- If you receive Idacio while you are pregnant, it is important that you inform your baby's doctor before your baby receives any vaccine. Your baby should not receive a 'live vaccine', such as BCG (used to prevent tuberculosis) within 5 months following your last Idacio injection during pregnancy.

During Idacio treatment

To make sure that Idacio is working properly and safely for you/your child, check in with your/your child's doctor regularly to discuss how you/your child are doing. Tell your/your child's doctor right away about any changes in your/your child's condition. Keep your/your child's doctor informed about how Idacio is working for you/your child.

- It is important to call your/your child's doctor right away about any unusual symptoms you/your child may have. This will help ensure you/your child get the right care. It will also lower the chance of a side effect becoming worse.
- Many side effects, including infections, can be managed if you tell your/your child's doctor right away.
- If you/your child get a side effect, your/your child's doctor will decide if you/your child should continue or stop your/your child's Idacio treatment. It is important to talk with your/your child's doctor to find out what is right for you/your child.
- Since side effects can happen after your/your child's last dose of Idacio, tell your/your child's doctor about any problems that you/your child may have had up to 70 days after your/your child's last injection of Idacio.
- Babies born to mothers who received Idacio during pregnancy should not receive any live vaccinations up until 5 months after the last dose of Idacio was given to the mother.

Tell your/your child's doctor about:

- Any new medical conditions that you/your child have;
- New medicines you/your child are taking;
- Any surgery or operation that you/your child have planned.

Some people taking Idacio may get serious side effects, including:

Infections
Idacio helps people with certain inflammatory diseases. It does this by blocking a part of the immune system. However, this part of the immune system also helps fight infection. This means Idacio can make