



## READ THIS

### INSPECT BAG PRIOR TO ACTIVATION

- SmofKabiven® is a three chamber bag
  - One chamber is **WHITE**
  - Two chambers are **CLEAR**
- **Discard bag if:**
  - Overwrap is **OPENED** or **DAMAGED**
  - More than one chamber is **WHITE**
  - Solution is **YELLOW**
  - Seals are already **BROKEN**

### REMOVE OVERWRAP

- Place bag on a clean, flat surface
- Tear from overwrap notch, located close to the ports (1)



- Tear long sides open to access the inner bag (2)



### ACTIVATE BAG

- Place bag on a clean, flat surface with text side up and ports pointing away from you
- Roll **tightly** from top of bag toward ports (3-4)



- **Apply pressure** until both vertical seals break and entire contents are white
- It may take up to 5 seconds of continued pressure to break vertical seals



**NOTE:** Both vertical seals must be broken from bends to ports. Upper section of vertical seals and horizontal seal may remain unbroken.

- After both vertical seals are broken, mix contents thoroughly by inverting the bag at least three times to ensure a homogenous mixture

### INSPECT BAG TO CONFIRM ACTIVATION

- An activated bag has both vertical seals broken from bends to ports and entire contents are white



### MAKE ADDITIONS (if prescribed)



**WARNING:** Ensure additives are compatible

- Immediately before injecting additives, break off the **WHITE** tab on the additive port with the arrow pointing toward the bag (7)
- Hold base of additive port horizontally, insert the needle through the center of the additive port's septum and inject the additives with known compatibility (8)



- Mix thoroughly after each addition

**NOTE:** The membrane of the additive port is sterile for first use. Use aseptic technique for subsequent additions. The septum can be pierced no more than 10 times with the recommended needle size 18 - 23G 1 1/2 inches (40mm).

### SPIKE AND HANG BAG

- Place the activated bag on a clean and flat surface
- Break off the **BLUE** tab on the infusion port with the arrow pointing away from the bag (9)



- Hold the base of the infusion port and insert the spike by rotating your wrist slightly (10)



- Hang the bag by the hole below the handle (11)
- For single use only. Discard unused portions.

**NOTE:** Use an infusion set with a 1.2 micron in-line filter. Use infusion sets (according to ISO number 8536-4) with an external spike diameter of 5.5 - 5.7mm





## READ THIS

### INSPECT BAG PRIOR TO ACTIVATION

- SmofKabiven® is a three chamber bag
  - One chamber is **WHITE**
  - Two chambers are **CLEAR**
- **Discard bag if:**
  - Overwrap is **OPENED** or **DAMAGED**
  - More than one chamber is **WHITE**
  - Solution is **YELLOW**
  - Seals are already **BROKEN**

### REMOVE OVERWRAP

- Place bag on a clean, flat surface
- Tear from overwrap notch, located close to the ports (1)



- Tear long sides open to access the inner bag (2)



### ACTIVATE BAG

- Place bag on a clean, flat surface with text side up and ports pointing away from you
- Roll **tightly** from top of bag toward ports (3-4)



- **Apply pressure** until both vertical seals break and entire contents are white
- It may take up to 5 seconds of continued pressure to break vertical seals



**NOTE:** Both vertical seals must be broken from bends to ports. Upper section of vertical seals and horizontal seal may remain unbroken.

- After both vertical seals are broken, mix contents thoroughly by inverting the bag at least three times to ensure a homogenous mixture

### INSPECT BAG TO CONFIRM ACTIVATION

- An activated bag has both vertical seals broken from bends to ports and entire contents are white



### MAKE ADDITIONS (if prescribed)



**WARNING:** Ensure additives are compatible

- Immediately before injecting additives, break off the **WHITE** tab on the additive port with the arrow pointing toward the bag (7)
- Hold base of additive port horizontally, insert the needle through the center of the additive port's septum and inject the additives with known compatibility (8)



- Mix thoroughly after each addition

**NOTE:** The membrane of the additive port is sterile for first use. Use aseptic technique for subsequent additions. The septum can be pierced no more than 10 times with the recommended needle size 18 - 23G 1 1/2 inches (40mm).

### SPIKE AND HANG BAG

- Place the activated bag on a clean and flat surface
- Break off the **BLUE** tab on the infusion port with the arrow pointing away from the bag (9)



- Hold the base of the infusion port and insert the spike by rotating your wrist slightly (10)



- Hang the bag by the hole below the handle (11)
- For single use only. Discard unused portions.

**NOTE:** Use an infusion set with a 1.2 micron in-line filter. Use infusion sets (according to ISO number 8536-4) with an external spike diameter of 5.5 - 5.7mm